



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Surveying the Down syndrome mouse model resource identifies critical regions responsible for chronic otitis media

Citation for published version:

Bhutta, MF, Cheeseman, MT, Herault, Y, Yu, YE & Brown, SDM 2013, 'Surveying the Down syndrome mouse model resource identifies critical regions responsible for chronic otitis media', *Mammalian Genome*, vol. E-pub 26 September. <https://doi.org/10.1007/s00335-013-9475-x>

Digital Object Identifier (DOI):

[10.1007/s00335-013-9475-x](https://doi.org/10.1007/s00335-013-9475-x)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Mammalian Genome

Publisher Rights Statement:

Copyright The Author(s) 2013

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Surveying the Down syndrome mouse model resource identifies critical regions responsible for chronic otitis media

Mahmood F. Bhutta · Michael T. Cheeseman ·
Yann Herault · Yuejin E. Yu · Steve D. M. Brown

Received: 21 February 2013 / Accepted: 15 August 2013 / Published online: 26 September 2013
© The Author(s) 2013. This article is published with open access at Springerlink.com

Abstract Chronic otitis media (OM) is common in Down syndrome (DS), but underlying aetiology is unclear. We analysed the entire available mouse resource of partial trisomy models of DS looking for histological evidence of chronic middle-ear inflammation. We found a highly penetrant OM in the Dp(16)1Yey mouse, which carries a complete trisomy of MMU16. No OM was found in the Dp(17)1Yey mouse or the Dp(10)1Yey mouse, suggesting disease loci are located only on MMU16. The Ts1Cje, Ts1RhR, Ts2Yah, and Ts65Dn trisomies and the trans-homosomal Tc1 mouse did not develop OM. On the basis of these findings, we propose a two-locus model for chronic middle-ear inflammation in DS, based upon

epistasis of the regions of HSA21 not in trisomy in the Tc1 mouse. We also conclude that environmental factors likely play an important role in disease onset.

Introduction

Down syndrome (DS) is the most common human chromosomal abnormality, occurring in 1/750 live births, and is due to partial or total trisomy HSA21. Variable clinical features include skull and midface malformation, cardiac anomalies, hypotonia, delayed growth, developmental disorders, thyroid disease, obesity, and also chronic otitis media (OM). Most children with DS develop OM with effusion (OME), which is often chronic (COME) and leads to a hearing loss that can compound learning difficulties (Libb et al. 1985; Marcell and Cohen 1992). A longitudinal study of 79 children with DS reported that OME was diagnosed in 93 % at age one, falling to 68 % by age 5 (Barr et al. 2011). Similar findings have been reported in other series (Brooks et al. 1972; Balkany et al. 1979; Maurizi et al. 1985; Dahle and McCollister 1986; Hassmann et al. 1998; Kattan et al. 2000; Mitchell et al. 2003; Park et al. 2011). COME may persist, and at least 10–20 % of adults with DS have a conductive hearing loss (Evenhuis et al. 1992; Van Buggenhout et al. 1999). Middle-ear inflammation may also be subclinical (Roizen et al. 1994). Several authorities advise routine and regular audiological screening of children with DS (American Academy of Pediatrics 2001; Downs Syndrome Medical Interest Group 2004).

Standard treatments for COME are not as effective in children with DS. Grommets (ventilation tubes) in this population may not significantly improve hearing, have little if any effect on disease recurrence, and are not

M. F. Bhutta (✉)
Nuffield Department of Surgical Sciences, John Radcliffe
Hospital, University of Oxford, Room 6607 Level 6,
Headley Way, Oxford OX3 9DU, UK
e-mail: m.bhutta@doctors.org.uk

M. F. Bhutta · M. T. Cheeseman · S. D. M. Brown
MRC Harwell, Harwell Science and Innovation Campus,
Oxfordshire OX11 0RD, UK

M. F. Bhutta
Department of Otolaryngology Head and Neck Surgery, John
Radcliffe Hospital, Level LG1 West Wing, Headley Way,
Oxford OX3 9DU, UK

Y. Herault
Institut de Génétique, Biologie Moléculaire et Cellulaire et
Institut, Clinique de la Souris, 1 rue Laurent Fries, Parc
d'Innovation, BP 10142, 67404 Illkirch-Graffenstaden Cedex,
France

Y. E. Yu
Department of Cancer Genetics, CGP L2-320, Roswell Park
Cancer Institute, Elm & Carlton Streets, Buffalo,
NY 14263, USA

infrequently complicated by infection (Selikowitz 1993; Iino et al. 1999; Shott et al. 2001). Some advocate hearing aids as the preferred treatment for COME in DS (National Institute for Health and Clinical Excellence 2008). An understanding of the pathogenic mechanisms leading to chronic middle-ear inflammation in DS offers potential for targeted and more effective therapy of COME in this population. It may also provide insights into mechanisms of inflammation in nonsyndromic COME, which is also highly heritable (Casselbrant et al. 1999). Specifically, trisomy in DS could mimic hypermorphic polymorphisms underlying nonsyndromic disease.

The aetiology of chronic middle-ear inflammation in DS is not understood. Children with DS are more prone to infection as a result of defective mucosal immunology (Kusters et al. 2009). This could explain the onset of OM, but the persistent nonsuppurative inflammation seen in COME is not thought to reflect persistent bacterial infection. Several authorities suggest that an anatomical anomaly of the Eustachian tube in DS contributes to nonresolving disease (Bluestone et al. 2005), but the only published study on Eustachian tube patency in DS did not demonstrate the tube to be obstructed (White et al. 1984).

DS is presumed to arise as a consequence of either an extra copy of protein-coding sequences that are dosage-sensitive or an extra copy of non-protein-coding sequences that are regulatory or otherwise functional (Antonarakis et al. 2004; Gardiner and Costa 2006; Megarbane et al. 2009; Tan et al. 2009; Wiseman et al. 2009). HSA21 is the smallest human autosome, with only around 300 genes, and so perhaps only a handful of genes play a role in the phenotypic manifestations of DS. Individuals with partial trisomy of HSA21 offer serendipitous clues to identify critical regions for DS phenotypes. Initial analyses of such individuals suggested that all or most phenotypic manifestations of DS were due to a short critical region (McCormick et al. 1989; Rahmani et al. 1989), but more recent analysis refutes this and suggests that loci determining the DS phenotype are scattered across HSA21 (Lyle et al. 2004; Korb et al. 2009). This analysis had not been undertaken specifically for the OM phenotype, but given the rarity of partial trisomy HSA21 and the lack of a large database of genotype-phenotype correlation for aneuploidies, such analysis is difficult at present.

Readouts of the transcriptome of HSA21 have also been used to define critical genes. Data from human (Ait Yahya-Graison et al. 2007) and mouse (Lyle et al. 2004) show that only around a third of the genes on HSA21 are transcribed at more than the theoretical 1.5-fold increase, suggesting that the majority of trisomic loci are compensated for by

negative feedback that modulates transcriptional activity or mRNA stability. Although this does appear to narrow the possibilities, the method is not entirely reliable. Data of the transcriptome from mouse and human differ, as do data from different tissue types (Lyle et al. 2004). In addition, transcription is but one component of gene expression (Schwanhauss et al. 2011), so small changes in transcription could have a large phenotypic effect at some loci, and large changes in transcription could have little or no phenotypic effect at other loci.

There has been increasing interest in using mouse models to unravel pathobiology in DS. The syntenic regions to HSA21 are scattered across the mouse genome on MMU16, MMU17, and MMU10 (see Fig. 1a), which makes genetic modelling difficult (Gardiner et al. 2003). However, there is a growing library of mouse models of DS. The Ts65Dn mouse (Davisson et al. 1993), Ts1Cje mouse (Sago et al. 1998), Ts2Yah mouse (Herault et al. 2009), Ts1Rhr mouse (Olson et al. 2004), Dp(16)1Yey mouse (Li et al. 2007), Dp(10)1Yey mouse (Yu et al. 2010), and Dp(17)1Yey mouse (Yu et al. 2010) each carry a partial trisomy for the homologous genes of HSA21. Together these models cover almost all regions triplicated in DS (Fig. 1b). In addition, the Tc1 mouse is a trans-chromosomal model, carrying most of HSA21 in addition to the full complement of mouse chromosomes (O'Doherty et al. 2005).


This mouse library has been surveyed to identify loci responsible for some of the phenotypic manifestations of DS, but it has never been systematically interrogated for the presence of OM. The Ts65Dn mouse, which carries a 15.6-Mb segmental trisomy of MMU16 (Duchon et al. 2011; Reinholdt et al. 2011), has been reported to develop spontaneous chronic OM (Han et al. 2009), and recently the Tc1 mouse has been reported to not develop chronic OM (Kuhn et al. 2012). There are no reports of the presence or absence of OM in the other murine models of DS. In this study we systematically screened the murine resource of DS models to identify the histological presence of chronic OM, with the aim of narrowing and defining a critical region or set of loci underlying this phenotype.

Materials and methods

Mouse strains

We analysed cadaveric specimens of almost the entire reported library of DS mouse models. Specimens of the Ts65Dn mouse, the Ts1Cje mouse, the Ts2Yah mouse, and the Ts1Rhr mouse were obtained from the Institut Clinique de la Souris, Strasbourg. Specimens of the Dp(16)1Yey

Fig. 1 Map of **a** murine synteny to human chromosome 21 and of **b** trisomic regions in mouse models of DS used in this study (showing genes at chromosome breakpoints). The syntenic regions to HSA21 are on MMU16 (76–98 Mb), MMU17 (31–32 Mb), and MMU10 (78–76 Mb). The Tc1 mouse carries an incomplete HSA21 in addition to the normal mouse genome, whereas all other models are trisomic for murine DNA. Known genes are derived from the Ensembl web browser



mouse, the Dp(10)1Yey mouse, and the Dp(17)1Yey mouse were obtained from Roswell Park, NY, USA. Specimens of the Tc1 mouse and additional specimens of the Ts65Dn mouse were obtained from University College London. The number and age of mice supplied depended upon local availability. All mice were supplied with wild-type (WT) littermate controls, and all were genotyped at source. The microbial status and housing conditions of colony mice varied between institutions, as detailed in Table 1.

Histology

Mouse heads were fixed in 10 % neutral buffered formalin, decalcified, and embedded in paraffin wax. Three-micron sections were stained with haematoxylin and eosin and assessed for histological evidence of inflammation of the middle ear (Cheeseman et al. 2011).

Statistics

We used Fisher's exact test to compare prevalence of OM in mutant versus WT littermates.

Results

We examined serial sections of mouse heads from eight mouse lines [Ts65Dn, Ts1Cje, Ts2Yah, Ts1Rhr, Dp(16)1Yey, Dp(10)1Yey, Dp(17)1Yey, and Tc1] for histological evidence of inflammation of the middle ear. A total of 532 ears were analysed and the results are documented in Table 2. For one mouse line, Ts65Dn, mice were gathered from two sources.

Only the Dp(16)1Yey mouse line reliably developed statistically significant chronic OM (19 ears in 10 of 13

Table 1 Microbial status of mouse strains

Source	Mouse lines	FELASA listed microbial agents	Housing conditions
Institut Clinique de la Souris	Ts65Dn, Ts1Cje, Ts2Yah, Ts1Rhr	Norovirus, <i>Helicobacter</i> spp., <i>Pseudomonas aeruginosa</i> , <i>Entamoeba muris</i>	Individually ventilated cages with HEPA-filtered air
Roswell Park	Dp(10)1Yey, Dp(16)1Yey, Dp(17)1Yey	Norovirus, <i>Helicobacter</i> spp.	Individually ventilated cages with HEPA-filtered air
University College London	Tc1, Ts65Dn	<i>Helicobacter hepaticus</i> , <i>Tritrichomonas</i>	Open cages

FELASA Federation for Laboratory Animal Science Associations

Table 2 Mouse strains used in this study, their source, and presence or absence of OM on histology (denominator is for number of ears)

Mouse line	Background strain	Trisomic segment	Source	Age range (months)	OM in mutant	OM in WT controls	<i>p</i> value
Dp(17)1Yey	B6;129	MMU17 <i>Abcg1-Rrp1b</i>	Roswell Park	2–3	0/28	1/32	1
Dp(10)1Yey	B6;129	MMU10 <i>Prmt2-Pdxk</i>	Roswell Park	2–4	0/28	0/28	1
Dp(16)1Yey	B6;129	MMU16 <i>Lipi-Znf295</i>	Roswell Park	2–15	19/26	7/24	0.004
Ts1Cje	B6	MMU16 <i>Sod1-Znf295</i>	ICS	2	2/28	0/28	0.491
Ts1Rhr	B6	MMU16 <i>Cbr1-Mx2</i>	ICS	2	0/36	0/36	1
Ts2Yah	B6	MMU16 <i>Stch-App</i>	ICS	2	0/34	0/34	1
Ts65Dn	C3H;B6	MMU16 <i>Mrpl39-Znf295</i> & MMU17 <i>Synj2-D17Mit58</i>	ICS	2	2/40 ^a	0/40	0.494
Ts65Dn	C3H;B6	MMU16 <i>Mrpl39-Znf295</i> & MMU17 <i>Synj2-D17Mit58</i>	UCL	11–16	1/12 ^b	0/22	0.353
Tc1	129;B6	HSA21 except <i>Rh46998-D21s11</i> & <i>Ifngr2-Rh120345</i>	UCL	2–3	0/30	0/26	1

The *p* values relate to Fisher's exact test for the difference in OM prevalence in mutants versus wild-type (WT) littermates

UCL University College London, ICS Institut Clinique de la Souris

^a Two ears in this cohort of Ts65Dn mice showed mild oedema of the middle-ear mucoperiosteum, but without leucocyte infiltration

^b One ear showed a small middle-ear exudate with micropolyp formation

mice). Histology revealed thickened polypoid mucoperiosteum with neutrophil and macrophage infiltration, and an effusion of variable cellularity (Fig. 2). In this line, disease was consistently present from the age of 2 months. Some of the older (>7 months old) WT littermate controls for this line also developed chronic OM (7 ears in 4 mice), but OM was not present in any of the younger WT littermates.

OM in other models was not present or it was rarely present. Given the reported OM in Ts65Dn mice, we obtained animals from two different sources. We found no evidence of OM in either colony. A unilateral small volume of exudate and inflammatory polyp formation were noted in one of the Ts65Dn mice from University College London (Fig. 2). There was also mild unilateral oedema in the middle-ear mucosa of two Ts65Dn mice from the Institut Clinique de la Souris, but without a leucocyte infiltrate (not shown). There was bilateral OM in one Ts1Cje mouse, with infiltration by macrophages and neutrophils (Fig. 2).

Discussion

Chromosomal localisation of OM

In this study we screened the DS mouse library to identify a critical region for genetic susceptibility to chronic OM. The Dp(16)1Yey mouse, which is trisomic for the entire region of MMU16 syntenic to HSA21, was found to reliably develop chronic OM. Here, OM was early onset (by 2 months) and with penetrance of 0.73 (19/26 ears). A few of the WT littermate controls of the Dp(16)1Yey mice also developed OM, but only when they were older than 7 months. Chronic OM is common in older laboratory mice (Haines et al. 2001), although the reasons are unclear, and so the pathology in the older WT mice is not unexpected.

Of note, there was no evidence of OM in the Dp(17)1Yey and Dp(10)1Yey models, suggesting that only genes found on MMU16 are responsible for the OM




Fig. 2 Histological section through the middle ear of a Dp(16)1Yey mouse, a Ts65Dn mouse, and a Ts1Cje mouse affected by chronic OM (H&E stain). There is polypoid hyperplasia of the mucoperosteum (m) with an effusion (e) infiltrated by neutrophils and macrophages. This was present in 19/26 of Dp(16)1Yey mice examined, 3/52 of Ts65Dn mice, and 1/14 Ts1Cje mice

phenotype. We also found no evidence of OM in the Tc1 mouse, in agreement with recently reported results (Kuhn et al. 2012). The Tc1 model carries many of the genes found in trisomy in DS (in the form of mouse and human chromosomes), but it has been reported to lack many elements of the DS phenotype. It has been surmised that this may reflect species differences in genetic regulatory elements, or it may be the result of genetic mosaicism (Patterson 2009). More recent work also suggests previously unrecognised gene rearrangements in the Tc1 mouse (E. Fisher, personal communication).

Our findings strongly implicate a region of MMU16 to underlie the chronic OM phenotype. Empirically, this region could be narrowed further by analysis of the

Table 3 Genes located in the regions *Rh46998-D21s11* (region A) and *Ifngr2-Rh123045* (region B)

Gene	Relative transcript level	Function	Candidate
Region A			
<i>CXADR</i>	?	Coxsackie and adenovirus receptor, maintains tight junctions	✖
<i>C21ORF</i>	?	Unknown	
<i>CHODL</i>	?	Endocytosis of glycoproteins and sugar-bearing pathogens	
<i>TMPRSS15</i>	?	Pancreatic enzyme	
Region B			
<i>IFNGR2</i>	1.37	Receptor subunit for interferon γ	✖
<i>TMEM50B</i>	1.43	Brain development	
<i>DNAJC28</i>	–	May act in protein folding or as a chaperone	
<i>GART</i>	1.46	Purine biosynthesis	
<i>SON</i>	1.77	mRNA splicing cofactor	
<i>DONSON</i>	1.35	Unknown	
<i>CRYZL1</i>	1.40	Unknown	
<i>ITSN1</i>	1.42	Coordinates endocytic membrane traffic	
<i>ATP5O</i>	1.40	ATP synthesis	
<i>MRPS6</i>	1.28	Mitochondrial protein synthesis	
<i>SLC5A3</i>	0.91	Prevents intracellular accumulation of myo-inositol	
<i>KCNE2</i>	1.32	Component of voltage-gated potassium channels	
<i>FAM165B</i>	1.39	Unknown	
<i>KCNE1</i>	–	Component of voltage-gated potassium channels	
<i>RCAN1</i>	0.79	Inhibits calcineurin-dependent signaling pathways	
<i>CLIC6</i>	–	Probable chloride ion channel	
<i>RUNX1</i>	1.31	Functions in haematopoiesis, interacts with TGF- β	✖

Regions A and B are trisomic in the Dp(16)1Yey mouse (develops OM), but not in the Tc1 mouse (no OM noted). Relative transcript levels for region B refer to those in adult lung of Ts65Dn mice (Lyle et al. 2004); region A is not trisomic in Ts65Dn mice. In the proposed multigenic model, genes from each of these regions interact to produce susceptibility to chronic OM. Highlighted “candidates” are those genes known to be involved in immune signaling

Ts65Dn, Ts1Cje, Ts1Rhr, and Ts2Yah mice, each of which carry shorter segments trisomic for MMU16 (Fig. 1).

Localisation of OM within MMU16

We found no OM in Ts1Rhr and Ts2Yah mice, but one Ts1Cje mouse developed severe bilateral OM. We observed definitive OM in only one ear of the 52 Ts65Dn ears screened, although there was mild mucosal oedema in two other Ts65Dn mouse ears (which may be indicative of early or mild inflammation). The absence of OM in these four mouse lines, coupled with the highly penetrant OM observed in the Dp(16)1Yey mouse, suggests a genetic model for OM in DS whereby more than one locus is required. Given the absence of OM in Tc1, this would suggest that the loci are present in the two chromosome 16 regions not found in Tc1, i.e., *Rh46998-D21s11* and *Ifngr2-Rh123045* (Fig. 1b). Known genes in these two regions and their function are detailed in Table 3. The first region contains 4 genes, of which only *Cxadr* is known to be involved in immune signalling. The second region contains 17 genes, of which two, *Ifngr2* and *Runx1*, are known to be involved in immune signalling. *Cxadr* and *Ifngr2* are usually invoked in response to viral rather than bacterial infection. Whether these or other loci in these regions are responsible for persistent middle-ear inflammation we can only surmise.

In considering this multigenic model, it is important to examine the differing results reported in this study and those of Han et al. (2009) regarding the incidence of OM observed in the Ts65Dn mouse. Han et al. reported a highly penetrant chronic OM in the Ts65Dn model (in 11/15 mice), with a variable cellular infiltrate and effusion, accompanied by audiological evidence of disease onset by the age of 2 months. The variable penetrance of OM in the Ts65Dn mouse observed in this and the Han study suggests that development of chronic OM in DS is not determined solely by a critical genetic region(s) in trisomy, but also by additional important modifiers of phenotype, possibly genetic, environmental, or both.

Background sequence variation could affect phenotype penetrance and expressivity. The Ts65Dn mice in our study were maintained on a C3;B6 background through breeding with local stock. The Ts65Dn mice reported by Han et al. were similarly maintained by local breeding on a C3;B6 background. However, even small genomic variations can play a pivotal role in phenotype penetrance in DS (Antonarakis et al. 2004) and thus interlaboratory variation might affect the OM phenotype. For example, allelic variation in *CRELD1* on HSA3 modifies the risk of cardiac malformation in children (Maslen et al. 2006). In the mouse, breeding the Ts1Rhr mouse model onto a mixed B6;C3H/129 background, as opposed to pure B6, leads to loss of the craniofacial phenotype (Keane et al. 2011). However, it is noteworthy that we obtained Ts65Dn mice from two separate sources, and in neither case did we observe OM.

An alternative, and perhaps more likely, explanation is that an environmental antigen, such as a bacterial pathogen, can

trigger the onset of OM in Ts65Dn mice. Variation in housing conditions could lead to differences in environmental exposure and so explain variable phenotype penetrance. Han et al. (2009) reported coagulase-negative *Staphylococci*, *Burkholderia cepacia*, and *Bordetella avium* in middle-ear effusions of their Ts65Dn mice, but also *Klebsiella oxytoca*, *Streptococcus viridans*, and *B. avium* in wild-type controls. We did not culture bacteria from effusions in this study (only fixed tissues were used), but because a number of microbes have been isolated at the source of these animals (Table 2), it is possible that one or more of these microbes, or other unidentified microbes, play a pivotal role in disease onset.

In summary, our data suggest a multigenic model for development of OM in DS mice, whereby loci from two chromosome 16 regions contribute to the onset of the disease. Consistent with our proposed model, we failed to observe definitive OM in the Ts65Dn mouse, which would carry only one of the required chromosome 16 regions. However, it may be that environmental insults or other genetic variation elsewhere in the genome may be sufficient along with the trisomic region present in Ts65Dn mouse to elicit OM. This would potentially account for the differences in incidence of OM observed between the work of Han et al. and the work reported here.

Acknowledgments M. Bhutta was funded by a Wellcome Trust Clinical Research Fellowship. Y. E. Yu was supported by the NIH (R01HL091519 and R01NS066072). We thank Frances Wiseman (UCL Institute of Neurology) for the gift of Ts65Dn tissue. We thank the pathology and histology teams at MRC Harwell for producing the histology sections for review.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Ait Yahya-Graison E, Aubert J et al (2007) Classification of human chromosome 21 gene-expression variations in Down syndrome: impact on disease phenotypes. *Am J Hum Genet* 81(3):475–491
- American Academy of Pediatrics (2001) Health supervision for children with Down syndrome. *Pediatrics* 107(2):442–449
- Antonarakis SE, Lyle R et al (2004) Chromosome 21 and Down syndrome: from genomics to pathophysiology. *Nat Rev Genet* 5(10):725–738
- Balkany TJ, Downs MP et al (1979) Hearing loss in Down's syndrome. A treatable handicap more common than generally recognized. *Clin Pediatr (Phila)* 18(2):116–118
- Barr E, Dungworth J et al (2011) The prevalence of ear, nose and throat disorders in preschool children with Down's syndrome in Glasgow. *Scott Med J* 56(2):98–103
- Bluestone CD, Bluestone MB, Coulter J (2005) The Eustachian tube: structure, function, role in otitis media. BC Decker, Hamilton
- Brooks DN, Wooley H et al (1972) Hearing loss and middle ear disorders in patients with Down's syndrome (mongolism). *J Ment Defic Res* 16(1):21–29

- Casselbrant ML, Mandel EM et al (1999) The heritability of otitis media: a twin and triplet study. *JAMA* 282(22):2125–2130
- Cheeseman MT, Tyrer HE et al (2011) HIF-VEGF pathways are critical for chronic otitis media in Junbo and Jeff mouse mutants. *PLoS Genet* 7(10):e1002336
- Dahle AJ, McCollister EP (1986) Hearing and otologic disorders in children with Down syndrome. *Am J Ment Defic* 90(6):636–642
- Davison MT, Schmidt C et al (1993) Segmental trisomy as a mouse model for Down syndrome. *Prog Clin Biol Res* 384:117–133
- Downs Syndrome Medical Interest Group (2004) Basic medical surveillance essentials for people with Down syndrome: hearing impairment. DSMIG, Nottingham
- Duchon A, Raveau M et al (2011) Identification of the translocation breakpoints in the Ts65Dn and Ts1Cje mouse lines: relevance for modelling Down syndrome. *Mamm Genome* 22(11–12):674–684
- Evenhuis HM, van Zanten GA et al (1992) Hearing loss in middle-age persons with Down syndrome. *Am J Ment Retard* 97(1):47–56
- Gardiner K, Costa AC (2006) The proteins of human chromosome 21. *Am J Med Genet C Semin Med Genet* 142C(3):196–205
- Gardiner K, Fortna A et al (2003) Mouse models of Down syndrome: how useful can they be? Comparison of the gene content of human chromosome 21 with orthologous mouse genomic regions. *Gene* 318:137–147
- Haines DC, Chattopadhyay S et al (2001) Pathology of aging B6;129 mice. *Toxicol Pathol* 29(6):653–661
- Han F, Yu H et al (2009) Otitis media in a mouse model for Down syndrome. *Int J Exp Pathol* 90(5):480–488
- Hassmann E, Skotnicka B et al (1998) Distortion products otoacoustic emissions in diagnosis of hearing loss in Down syndrome. *Int J Pediatr Otorhinolaryngol* 45(3):199–206
- Herauld Y, Lopes Pereira P et al (2009) Tackling the complexity of the genotype–phenotype relationship in the Down syndrome with the mouse aneuploidy zoo: a resource of new models to study aneuploidies involving human chromosome 21. The American Society of Human Genetics 59th Annual Meeting, Honolulu HI, 20–24 October 2009
- Iino Y, Imamura Y et al (1999) Efficacy of tympanostomy tube insertion for otitis media with effusion in children with Down syndrome. *Int J Pediatr Otorhinolaryngol* 49(2):143–149
- Kattan HA, Jarrar RF et al (2000) A pilot study of the relationship between Down's syndrome and hearing loss. *Saudi Med J* 21(10):931–933
- Keane TM, Goodstadt L et al (2011) Mouse genomic variation and its effect on phenotypes and gene regulation. *Nature* 477(7364):289–294
- Korbel J, Tirosh-Wagner T et al (2009) The genetic architecture of Down syndrome phenotypes revealed by high-resolution analysis of human segmental trisomies. *Proc Natl Acad Sci USA* 106(29):12031–12036
- Kuhn S, Ingham N et al (2012) Auditory function in the tc1 mouse model of Down syndrome suggests a limited region of human chromosome 21 involved in otitis media. *PLoS One* 7(2):e31433
- Kusters MA, Verstegen RH et al (2009) Intrinsic defect of the immune system in children with Down syndrome: a review. *Clin Exp Immunol* 156(2):189–193
- Li Z, Yu T et al (2007) Duplication of the entire 22.9 Mb human chromosome 21 syntenic region on mouse chromosome 16 causes cardiovascular and gastrointestinal abnormalities. *Hum Mol Genet* 16(11):1359–1366
- Libb JW, Dahle A et al (1985) Hearing disorder and cognitive function of individuals with Down syndrome. *Am J Ment Defic* 90(3):353–356
- Lyle R, Gehrig C et al (2004) Gene expression from the aneuploid chromosome in a trisomy mouse model of Down syndrome. *Genome Res* 14(7):1268–1274
- Marcell MM, Cohen S (1992) Hearing abilities of Down syndrome and other mentally handicapped adolescents. *Res Dev Disabil* 13(6):533–551
- Maslen CL, Babcock D et al (2006) CRELD1 mutations contribute to the occurrence of cardiac atrioventricular septal defects in Down syndrome. *Am J Med Genet A* 140(22):2501–2505
- Maurizi M, Ottaviani F et al (1985) Audiological findings in Down's children. *Int J Pediatr Otorhinolaryngol* 9(3):227–232
- McCormick MK, Schinzel A et al (1989) Molecular genetic approach to the characterization of the Down syndrome region of chromosome 21. *Genomics* 5(2):325–331
- Megarbane A, Ravel A et al (2009) The 50th anniversary of the discovery of trisomy 21: the past, present, and future of research and treatment of Down syndrome. *Genet Med* 11(9):611–616
- Mitchell RB, Call E et al (2003) Ear, nose and throat disorders in children with Down syndrome. *Laryngoscope* 113(2):259–263
- National Institute for Health and Clinical Excellence (2008) Surgical management of otitis media with effusion in children. NICE, London
- O'Doherty A, Ruf S et al (2005) An aneuploid mouse strain carrying human chromosome 21 with Down syndrome phenotypes. *Science* 309(5743):2033–2037
- Olson LE, Richtsmeier JT et al (2004) A chromosome 21 critical region does not cause specific Down syndrome phenotypes. *Science* 306(5696):687–690
- Park AH, Wilson MA et al (2011) Identification of hearing loss in pediatric patients with Down syndrome. *Otolaryngol Head Neck Surg* 146(1):135–140
- Patterson D (2009) Molecular genetic analysis of Down syndrome. *Hum Genet* 126(1):195–214
- Rahmani Z, Blouin JL et al (1989) Critical role of the D21S55 region on chromosome 21 in the pathogenesis of Down syndrome. *Proc Natl Acad Sci USA* 86(15):5958–5962
- Reinholdt LG, Ding Y et al (2011) Molecular characterization of the translocation breakpoints in the Down syndrome mouse model Ts65Dn. *Mamm Genome* 22(11–12):685–691
- Roizen NJ, Martich V et al (1994) Sclerosis of the mastoid air cells as an indicator of undiagnosed otitis media in children with Down's syndrome. *Clin Pediatr (Phila)* 33(7):439–443
- Sago H, Carlson EJ et al (1998) Ts1Cje, a partial trisomy 16 mouse model for Down syndrome, exhibits learning and behavioral abnormalities. *Proc Natl Acad Sci USA* 95(11):6256–6261
- Schwanhauser B, Busse D et al (2011) Global quantification of mammalian gene expression control. *Nature* 473(7347):337–342
- Selikowitz M (1993) Short-term efficacy of tympanostomy tubes for secretory otitis media in children with Down syndrome. *Dev Med Child Neurol* 35(6):511–515
- Shott SR, Joseph A et al (2001) Hearing loss in children with Down syndrome. *Int J Pediatr Otorhinolaryngol* 61(3):199–205
- Tan TY, Aftimos S et al (2009) Phenotypic expansion and further characterisation of the 17q21.31 microdeletion syndrome. *J Med Genet* 46(7):480–489
- Van Buggenhout GJ, Trommelen JC et al (1999) Down syndrome in a population of elderly mentally retarded patients: genetic-diagnostic survey and implications for medical care. *Am J Med Genet* 85(4):376–384
- White BL, Doyle WJ et al (1984) Eustachian tube function in infants and children with Down's syndrome. In: Lim DJ, Bluestone CD, Klein JO, Nelson JD (eds), Recent advances in otitis media with effusion. Proceedings of the Third International Symposium, Fort Lauderdale, FL, May 17–20, 1983. Philadelphia: B C Decker, pp 62–66
- Wiseman FK, Alford KA et al (2009) Down syndrome: recent progress and future prospects. *Hum Mol Genet* 18(R1):R75–R83
- Yu T, Li Z et al (2010) A mouse model of Down syndrome trisomic for all human chromosome 21 syntenic regions. *Hum Mol Genet* 19(14):2780–2791